

Name \_\_\_\_\_

Student number \_\_\_\_\_

## SE 2A04 Fall 2001

### Surprise Quiz 1 Answer Key

Instructor: W. M. Farmer

Revised: 5 November 2001

You have 20 minutes to complete this quiz consisting of 3 pages and 10 questions. You may use your notes and textbooks. Circle the *best* answer for the multiple choice questions. Recall that the  $\supset$  symbol denotes implication. Each question is worth 10 points. Good luck!

(1) Let  $a$  denote an integer. What is the value of

$$\text{if}(a > 2 \supset a > 0, 17, 19) \text{ ?}$$

- (a) 2.
- (b) 0.
- (c) 17.
- (d) 19.

(2) Let  $r$  denote a real number. What is the value of

$$(\lambda x : \mathbf{R} . x^2)(r) \text{ ?}$$

- (a)  $(\lambda x : \mathbf{R} . r^2)$ .
- (b)  $(\lambda r : \mathbf{R} . r^2)$ .
- (c)  $r^2$ .
- (d)  $x^2$ .

(3) Let **Push**, **Top**, **Pop** be members of the Stacks interface (as given in class);  $i, j$  be of type **Integer**; and  $s$  be of type **Stack**. What is the value of

$$\text{Top}(\text{Push}(j, \text{Pop}(\text{Push}(i, s)))) \text{ ?}$$

- (a) **Push**.
- (b)  $s$ .
- (c)  $i$ .
- (d)  $j$ .

Test continues on next page.

1/3

(4) Let  $x$  and  $y$  denote real numbers. Which expression denotes the minimum of  $x$  and  $y$ ?

- (a)  $-\max(x, y)$ .
- (b)  $\text{if}(x \leq y, y, x)$ .
- (c)  $\text{if}(x \geq y, x, y)$ .
- (d)  $\boxed{\text{if}(x \leq y, x, y)}$ .

(5) Let  $a$  be a real number and  $S$  be a set of real numbers. Which statement says that  $a$  is an upper bound of  $S$ ?

- (a)  $\forall x : \mathbf{R} . x \leq a$ .
- (b)  $\boxed{\forall x : \mathbf{R} . x \in S \supset x \leq a}$ .
- (c)  $\exists x : \mathbf{R} . x \in S \wedge x \leq a$ .
- (d)  $\exists x : \mathbf{R} . x \in S \wedge x < a$ .

(6) Which statement says that **Push** is a one-to-one function?

- (a)  $\forall i_1, i_2 : \text{INTEGER}, s_1, s_2 : \text{Stack} .$   
 $\boxed{\text{Push}(i_1, s_1) = \text{Push}(i_2, s_2) \supset (i_1 = i_2 \wedge s_1 = s_2)}$ .
- (b)  $\forall i_1, i_2 : \text{INTEGER}, s_1, s_2 : \text{Stack} .$   
 $(i_1 = i_2 \wedge s_1 = s_2) \supset \text{Push}(i_1, s_1) = \text{Push}(i_2, s_2)$ .
- (c)  $\forall t : \text{Stack} . \exists i : \text{INTEGER}, s : \text{Stack} . t = \text{Push}(i, s)$ .
- (d)  $\exists i : \text{INTEGER}, s : \text{Stack} . \text{Push}(i, s)$ .

(7) Which relation describes a function from the integers to the real numbers?

- (a)  $\{(x, y) \in \mathbf{Z} \times \mathbf{R} \mid x \leq y\}$ .
- (b)  $\{(x, y) \in \mathbf{R} \times \mathbf{R} \mid y = x * x\}$ .
- (c)  $\boxed{\{(x, y) \in \mathbf{Z} \times \mathbf{R} \mid y = x + 17\}}$ .
- (d)  $\{(x, y) \in \mathbf{Z} \times \mathbf{R} \mid 19 = 19\}$ .

(8) Let  $\mathbb{N}$  denote the natural numbers. Which statement defines

$$f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$$

by recursion to be natural number multiplication?

- (a)  $\forall x, y : \mathbb{N} . f(x, y) = f(x - 1, y + 1)$ .
- (b)  $f = \lambda x, y : \mathbb{N} . \text{if}(x = 0, x, y)$ .
- (c)  $\boxed{\forall x, y : \mathbb{N} . f(x, y) = \text{if}(x = 0, 0, y + f(x - 1, y))}$ .
- (d)  $\forall x, y : \mathbb{N} . x * y$ .

(9) Let  $m$  be an Oberon array of type **REAL** indexed from 0 to 999. Which statement says exactly that the members of  $m$  indexed from 17 to 313 (inclusive) are ordered from greatest to least?

(a) 
$$\forall i, j : \text{INTEGER} . (17 \leq i \wedge i \leq 313 \wedge 17 \leq j \wedge j \leq 313 \wedge i \leq j) \supset m[i] \geq m[j].$$

(b)  $\forall i : \text{INTEGER} . (17 \leq i \wedge i \leq 313) \supset m[i] \geq m[i + 1].$

(c)  $\forall i : \text{INTEGER} . m[i] \geq m[i + 1].$

(d)  $\forall i : \text{INTEGER} . m[i - 1] \geq m[i + 1].$

(10) Which expression denotes the cubic root of 139?

(a)  $139 * 139 * 139.$

(b)  $\forall x, y : \mathbf{R} . x < 139^3 < y.$

(c)  $\exists x : \mathbf{R} . x * x * x = 139.$

(d) 
$$\boxed{\text{I} x : \mathbf{R} . x * x * x = 139.}$$